

Natural and Cultural History of the Golfo Dulce Region, Costa Rica

Anton WEISSENHOFER, Werner HUBER,
Veronika MAYER, Susanne PAMPERL, Anton WEBER,
Gerhard AUBRECHT (scientific editors)

HPSG.: Biologiezentrum der Oberösterreichischen Landesmuseen



Natural and Cultural History of the Golfo Dulce Region, Costa Rica

Historia natural y cultural de la
región del Golfo Dulce, Costa Rica

Anton WEISSENHOFER, Werner HUBER, Veronika MAYER, Susanne PAMPERL,
Anton WEBER, Gerhard AUBRECHT (scientific editors)



Impressum

- Katalog / Publication: Stapfia **88**, zugleich Kataloge der Oberösterreichischen Landesmuseen N.S. **80**
ISSN: 0252-192X
ISBN: 978-3-85474-195-4
- Erscheinungsdatum / Date of delivery: 9. Oktober 2008
- Medieninhaber und Herausgeber / Copyright: Land Oberösterreich, Oberösterreichische Landesmuseen, Museumstr.14, A-4020 Linz
Direktion: Mag. Dr. Peter Assmann
Leitung Biologiezentrum: Dr. Gerhard Aubrecht
Url: <http://www.biologiezentrum.at>
E-Mail: bio-linz@landesmuseum.at
In Kooperation mit dem Verein zur Förderung der Tropenstation La Gamba (www.lagamba.at).
- Wissenschaftliche Redaktion / Scientific editors: Anton Weissenhofer, Werner Huber, Veronika Mayer, Susanne Pamperl, Anton Weber, Gerhard Aubrecht
Redaktionsassistent / Assistant editor: Fritz Gusenleitner
Layout, Druckorganisation /
Layout, printing organisation: Eva Rührnöbl
Druck / Printing: Plöchl-Druck, Werndlstraße 2, 4240 Freistadt, Austria
Bestellung / Ordering: <http://www.biologiezentrum.at/biophp/de/stapfia.php> oder / or bio.buch@landesmuseum.at
Das Werk einschließlich aller seiner Teile ist urheberrechtlich geschützt. Jede Verwertung außerhalb der engen Grenzen des Urheberrechtsgesetzes ist ohne Zustimmung des Medieninhabers unzulässig und strafbar. Das gilt insbesondere für Vervielfältigungen, Übersetzungen, Mikroverfilmungen sowie die Einspeicherung und Verarbeitung in elektronischen Systemen. Für den Inhalt der Abhandlungen sind die Verfasser verantwortlich. Schriftentausch erwünscht!
All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means without prior permission from the publisher. We are interested in an exchange of publications.
- Umschlagfoto / Cover: Blattschneiderameisen. Photo: Alexander Schneider. Layout: E. Rührnöbl.
- Zitiervorschlag für das Buch / The complete book may be referenced as follows: Weissenhofer, A., Huber W., Mayer V., Pamperl S., Weber A. & G. Aubrecht (Hrsg.; 2008): Natural and cultural history of the Golfo Dulce region, Costa Rica. — Stapfia 88: 768 pp.
- Zitiervorschlag für Einzelarbeiten / Single contributions may be referenced as follows: Weissenhofer A., Huber W. & M. Klingler (2008): Geography of the Golfo Dulce region. — Stapfia 88: #-#.
- Ausstellung / Exhibition: Der Pfad des Jaguars. Tropenstation La Gamba, Costa Rica
Ort / Address: Biologiezentrum der Oberösterreichischen Landesmuseen, J.-W.-Klein-Straße 73, 4040 Linz, Austria
Zeitraum / Period: 10. Oktober 2008 bis 22. März 2009
- Konzept, Organisation und Gestaltung /
Concept, organization, design: Mag. Stephan Weigl, Mag. Dr. Werner Huber, Mag. Dr. Anton Weissenhofer, Daniel Schaber
Ausstellungstechnik, Mitarbeit /
Exhibition techniques, collaboration: Jürgen Plass, Roland Rupp, Bruno Tumfart, Erwin Kapl, Josef Schmidt, Roland Zarre, Michaela Minich, Charlotte Füreder, Georg Proske, Franz Rammerstorfer
- Museumspädagogik / Museum education: Mag. Claudia Kiesenhofer
Leihgeber / Lenders: Naturhistorisches Museum Wien, Mag. Dr. Werner Huber, Mag. Dr. Anton Weissenhofer, Universität Wien; Mag. Felix Holzmann, Bischofshofen

Due to the orographic formation of its interior and its humid climate, the Golfo Dulce Region is rich with biodiversity, containing very dense flora and fauna. After HOLDRIDGE (1971), the region was subdivided into different zones, including the tropical rainforest, the tropical wetland forest, and tropical premontane rainforest. The biogeographical situation in this area shows many similarities to the flora and fauna in the Amazon and the Colombian Chocó Region and serves as a land bridge with a valuable genetic base between North and South America. After unregulated seizure of land by agricultural settlers, lumberjacks, and large landowners in the 1940s and 1950s, regulated, state-subsidised settlement reform intended to support agricultural exports in the 1960s, and intensification of the livestock industry in the 1970s, primary and secondary forest reserves have shrunk to a minimum. The constant expansion of monocultures on new land has far-reaching consequences for the local ecosystem.

The conservation and sustainable use of tropical forests is established in the Forest Declaration, Convention on Climate Protection, and Convention on the Protection of Species, which demonstrate worldwide concern for these issues. As a regional example, in the 4,304.80 km² drainage basin, the ACOSA (Área de Conservación OSA), which covers an area spanning the Cantons Osa, Golfito und Corredores, aims to protect species diversity within the 17 game preserves, which are 44.7% covered by forest, through integration and an alliance with the Parques Nacionales, Vida Silvestres y Forestales (Fig. 2). The main sector of the Corcovado National Park on the Osa Peninsula covers 424 km² and the Piedras Blancas National Park covers 148 km². The altitude ranges from sea level to 745 m on the Osa Peninsula (Cerro Rincón and Cerro Mueller in the Fila Matajambre) and to 579 m in the Esquinas forest (Cerro Nicuesa). The Golfo Dulce Forest Reserve (592 km²) was established between the two parks, thereby forming a natural forest corridor.

References

- CHINCHILLA V.E. (1987): Atlas cantonal de Costa Rica. — IFAM. San José.
- HARTSHORN, G.S. (1983): Plants: introduction. — In: JANZEN D.H. (ed.), Costa Rican natural history. Chicago: Univ. Chicago Press: 118-157.
- HERRERA-MACBRYDE O., MALDONADO V. JIMÉNEZ T.R. & K. THOMSEN (1997): Osa Peninsula and Corcovado National Park, Costa Rica. — In: DAVIS S.D., HEYWOOD V.H., HERRERA-MACBRYDE O., VILLA-LOBOS J. & A.C. HAMILTON (eds), Centres of plant diversity. A guide and strategy for their conservation. Vol. 3. WWF, IUCN.
- HERWITZ S.R. (1981): Regeneration of selected tropical tree species in Corcovado National Park, Costa Rica. — Univ. Calif. Publ. Geogr. 24.
- HOLDRIDGE L.R. (1971): Forest environments in tropical life zones. A pilot study. — Oxford.
- MALZER O. (2001): Geological History of Central America and the Golfo Dulce Region. — *Stapfia* 78: 34-46.
- MORA C.S. (1990): La Geología y sus procesos. — San José.
- TOSI J.A. Jr. (1975): The Corcovado Basin on the Osa Peninsula. — In: Tosi J.A. Jr. (ed.), Potential national parks, nature reserves, and wildlife sanctuary areas in Costa Rica: a survey of priorities. San José: Centro Científico Tropical. Separate pp. 12.

Addresses of authors:

Anton WEISSENHOFER
Werner HUBER

Department of Palynology and Structural Botany
Faculty Center of Botany
University of Vienna
Rennweg 14
A-1030 Vienna, Austria

E-mail: anton.weissenhofer@univie.ac.at
werner.huber@univie.ac.at

Michael KLINGLER
Workgroup Development Studies
and Sustainability Science
Institute of Geography
University of Innsbruck
Innrain 52

A-6020 Innsbruck, Austria
E-mail: michael.klingler@uibk.ac.at

Plant diversity and biogeography of the Golfo Dulce region, Costa Rica

Diversidad vegetal y biogeografía de la región de Golfo Dulce, Costa Rica

Werner HUBER, Anton WEISSENHOFER, Nelson ZAMORA & Anton WEBER

Abstract: With about 2.700 species of vascular plants, the Golfo Dulce region is one of the most floristically diverse regions of Central America. After a brief historical survey of botanical work in Costa Rica and in the Golfo Dulce region, the paper addresses recent investigations on tree diversity and discusses the reasons for the great botanical richness of the region. A biogeographical analysis of the tree species of the Esquinas forest reveals a close relation to northern South America, especially the Chocó region (parts of Colombia, Ecuador and Peru). The high degree of plant endemism in the Golfo Dulce region is remarkable. An analysis of seven representative families shows that about 18% of the investigated species are endemic. In the herbaceous Marantaceae, even 32% of the species are endemic. The reasons assumed responsible are addressed.

Key words: Costa Rica, Golfo Dulce, flora, biodiversity, biogeography, plant diversity, endemism.

Resumen: Con aproximadamente 2.700 especies de plantas vasculares, la región de Golfo Dulce es una de las áreas vegetacionales más diversas de América Central. Después de una breve investigación histórica sobre el trabajo botánico en Costa Rica y en la región de Golfo Dulce, este documento aborda las recientes investigaciones sobre la diversidad arbórea y discute las razones de la elevada riqueza vegetal de la región. Un análisis biogeográfico de las especies arbóreas del bosque Esquinas, revela una estrecha relación con el norte de América del Sur, especialmente la región de Chocó (parte de Colombia, Ecuador y Perú). Se destaca el alto grado de plantas endémicas en la región de Golfo Dulce. Un análisis de siete familias representativas, muestran que aproximadamente el 18% de las especies estudiadas son endémicas. La familia Marantaceae presenta un 32% de especies endémicas.

Palabras clave: Costa Rica, Golfo Dulce, flora, biodiversidad, biogeografía, diversidad de plantas, endemismo.

Introduction

The present paper presents a brief survey of plant (especially tree) diversity in the Golfo Dulce area and the plants' biogeographical relations. Starting from a short overview on the botanical work carried out in Costa Rica in the 19th and 20th centuries, recent and current floristic studies and biogeographical analyses are considered. The reasons for the close affinities to the flora of north-western South America and the noticeable endemism are addressed.

Floristic work in Costa Rica – a brief history

Out of all the Central American countries, Costa Rica has received the greatest attention from botanical collectors. While most parts of Central America were explored by Spanish botanists in colonial times, this was not so in Costa Rica. Botanists from various Euro-

pean countries worked in Costa Rica in the 19th century (see GÓMEZ 1986 and GRAYUM et al. 2004). Among the most prominent were Anders Sandøe Ørsted (1816-1872) from Denmark, Jozef v. Warscewicz (1812-1866) from Lithuania, Karl Hoffmann (1823-1859) from Germany, and Henri Pittier (1857-1950) and Adolphe Tonduz (1862-1921) from Switzerland. Names of many Costa Rican plant genera and species commemorate these scientists, e.g., *Oerstedella*, *Dieffenbachia oerstedii*, *Inga oerstediana*, *Hoffmannia*, *Warscewiczia*, *Cryosophila warscewiczii*, *Annona pittieri*, *Dracontium pittieri*, *Aristolochia tonduzii*, *Ficus tonduzii*, and many others. Significant contributions were also made by local collectors such as Anastasio Alfaro (1865-1951; *Alfaroa!*) and Juan José Cooper (1843-1911; *Ruellia cooperi*). Tonduz collected more than 18.000 specimens in Costa Rica. DURAND & PITTIER (1891-1901) were the first to publish a Costa Rican flora, the "Primitiae florae costaricensis", which included some 5.000 species. At the end of the 19th century, Central




Fig. 1: The ten most species-rich plant families (angiosperms) in the ACOSA region.

America and Costa Rica became politically attractive to the United States, and U.S. botanists started collection work in Costa Rica on a larger scale. John Donnell Smith (1829-1928), Paul C. Standley (1884-1963), Paul H. Allen (1911-1963) and William C. Burger (born 1932), have been among the most active botanists working in Costa Rica in the first half of the 20th century. Paul Standley collected more than 15,000 specimens and produced a four volume flora, the “Flora of Costa Rica” (STANDLEY 1937-1940). This included 6,085 species of higher plants. Standley recognised the country’s enormous plant diversity, stating: “Of all Central American countries Costa Rica possesses by far the richest flora”. According to GENTRY (1978), William Burger collected about 70,000 specimens for a new flora, the “Flora Costaricensis” (start of publication 1971). In a special paper, he discussed at length the question “Why are there so many kinds of flowering plants in Costa Rica?” (BURGER 1985). In 1985, Gomez-Laurito & Fournier counted about 860 native genera (in 192 families) of woody plants in Costa Rica. Knowledge of the Costa Rican flora has increased rapidly during recent decades. In a joint effort started in 1989, the Mis-




Fig. 2: The 15 most species-rich genera (angiosperms) in the ACOSA region.

souri Botanical Garden (USA), the Instituto Nacional de Biodiversidad (INBio, CR) and the Museo Nacional de Costa Rica (CR) published the first parts of the “Manual de Plantas de Costa Rica” (I-IV; HAMMEL et al. 2003a,b, 2004, 2007), covering 4,393 species, 817 genera and 104 families. At present, 9,361 species, 2,023 genera and 255 families of higher plants are known in Costa Rica (ZAMORA et al. 2004).

The botanical work that has been carried out around the Golfo Dulce must be considered in particular depth. Among the first botanists collecting extensively in the region was the Austrian botanist Georg Cufodontis (1896-1974) (see papers of DÍAZ and WEBER, this volume). In the 1950s, the US botanist Paul Allen, who lived in Palmar Sur, published the “The Rain Forests of Golfo Dulce” (ALLEN 1956), which remains a classic. At the end of the 1980s, INBio started to collect plants and to establish forest plots on the Osa Peninsula. The great amount of data is largely unpublished so far. Only a semi-popular field guide to the trees of the Osa Peninsula has appeared (QUESADA et al. 1997). In 1993, HUBER (1996a, 1996b, 2005) and WEISSENHOFER (1996, 1997, 2005) started botanical investigations in the Esquinas Forest (now Piedras Blancas National Park), establishing plots which have been monitored continuously ever since. In 2001 the “Introductory field guide to the flowering plants of the Golfo Dulce Rainforests, Costa Rica” (WEBER et al. 2001) was published.

Flora

The Golfo Dulce region is one of those rare places on earth where an enormous biological diversity can be found within a relatively small geographical area (VAUGHAN 1981). By any measure, it is one of the botanically most diverse regions of Central America. This is partly because of Costa Rica’s role as a “corridor” which enables migration of plants and animals from North to South America and vice versa. In 2001, 2,369 species of vascular plants – roughly a quarter of all Costa Rican plant species – were recorded from the Golfo Dulce area (WEISSENHOFER et al. 2001). At present, 2,662 species are recorded in the INBio database for the ACOSA region (see map, Fig. 3). At the family level, the flora contains 23 families of pteridophytes, two families of gymnosperms (with one species each), and 162 families of angiosperms. It is also notable that the region harbours over 700 tree species. This is the greatest tree species diversity known from Central America, and represents one quarter of all tree species found in Costa Rica (QUESADA et al. 1997).

The families with the highest species richness in the ACOSA region are shown in Fig. 1. The two largest families are Fabaceae (sensu lato; 193 spp.) and Rubia-

ceae (181 spp.), numbering almost 200 species each, followed by Orchidaceae and Melastomataceae, each with more than 100 species. These are followed by Araceae (93), Piperaceae (78), Poaceae (71), Euphorbiaceae (63), Asteraceae (58) Lauraceae (53), Moraceae (49), Solanaceae (45) and Arecaceae (44).

With regard to the generic level, the most diverse genera are shown in Fig. 2. The most speciose genera are *Piper* (Piperaceae) and *Psychotria* (Rubiaceae), with nearly 60 and 50 species, respectively.

In the recent past, about 62 species from the Golfo Dulce area have been described as new to science. Significant and conspicuous new species include *Porina pili-fera* (Lichenes), *Costus osaensis* (Costaceae), *Ruptiliocarpus caracolito* (Lepidobotryaceae), *Justicia peninsularis* (Acanthaceae), *Lycaste bruncana* (Orchidaceae), *Pleurothyrium golfodulcensis* (Lauraceae), *Aiouea obscura* (Lauraceae), *Ocotea patula* (Lauraceae), *Inga golfodulcensis* (Fabaceae/Mimosoideae), and *Stemmadenia paulii* (Apocynaceae). More than 50 species were recorded as new for the flora of Costa Rica, including *Ziziphus chloroxylon* (Rhamnaceae), *Oecopetalum greenmanii* (Icacinaceae), *Recchia simplicifolia* (Simaroubaceae), *Microrhophis venulosa* (Sapotaceae) and *Buchenavia tetraphylla* (Combretaceae). This shows that knowledge of plant diversity is still increasing and, more generally, that the tropical flora is badly known and under-collected (WEISSENHOFER et al. 2001). Ironically, most of the new discoveries were made along paths or near clearings.

Plant diversity and reasons assumed responsible

One of the most distinctive features of tropical forests worldwide, and neotropical lowland forests in particular, is the high diversity of plants and animals. The species richness tends to be highest in the wet lowlands, including the Golfo Dulce region. The Golfo Dulce area has an important climatic peculiarity; it experiences 1-3 months of low precipitation (WEISSENHOFER & HUBER, this volume). The slight seasonality seems to have a significant effect on species diversity. Moreover, there are special climatic conditions related to the position of the area: this is protected from north winds and drought effects by the presence and position of the Cordillera Talamanca. Rainfall is mostly due to southern pacific winds (HERRERA 1985). In addition, the area is geologically and geomorphologically unique with regard to its tectonic origin, hilly terrain, and peninsular situation, with the Osa Peninsula almost entirely surrounded by the sea.

There are other important reasons for the high diversity of plants. The region shows a very wide range of

different habitats, with natural habitats ranging from lowland rainforests to lower montane forest, and the inclusion of azonal vegetation such as mangroves, swamps, ponds, riverine vegetation etc.

As to altitude, three principal vegetation patterns can be distinguished:

(1) **Vegetation on alluvial soils.** This is mainly found in flat areas near or along the coast (Osa Peninsula) and includes the mangroves. The temperatures are distinctly higher than in the other types. Several floristic elements are restricted to this area.

(2) **Vegetation on hilly terrain,** located mainly between 40 and 500 m above sea level (a.s.l.). This covers most of the area, including the whole Esquinas forest. Here the highest species diversity is found.

(3) **Vegetation over 500 m a.s.l.** (up to 745 m a.s.l.: Cerro Rincon). This area experiences cloudy conditions. The occurrence of montane elements is characteristic, some belonging to temperate families and genera (Fagaceae: *Quercus insignis*, *Q. rapurahuensis*; Juglandaceae: *Oreomunnea pterocarpa*, *Alfaroa guanacastensis*; see ZAMORA et al. 2004 and SINAC 2007).

Generally, the forests around the Golfo Dulce have a strong spatial structure at the landscape scale and thus provide a great variety of ecosystems, microhabitats and ecological niches (see WEISSENHOFER et al., this volume). Rapid forest dynamics contribute to the maintenance of high diversity on a geographical gradient. Furthermore, the high edaphic heterogeneity and diversity of soils support a high plant-diversity (CONDIT et al. 2002). PAMPERL (2001) documented a high degree of soil heterogeneity in the research plots in the Esquinas forest. Today, man-made habitats such as pastures and agriculture land, found at all altitude levels, play a significant role.

A recent study (CASTILLO et al. 2007), analysing 76 one-hectare plots in order to determine the distribution and abundance of canopy trees species in the Osa Peninsula, reached the conclusion that in the western Golfo Dulce area five centres of tree diversity can be distinguished: (1) Mogos, (2) Bahía de Chal-Alto San Juan, (2) Rincón-Rancho Quemado, (3) La Palma, (4) Fila Carbonera, and (5) Corcovado National Park. The study recorded 5.227 individuals of 65 selected tree species (≥ 5 cm diameter at breast height) in the whole area. Among the most abundant species were *Symphonia globulifera* (759 individuals), *Carapa guianensis* (625), *Brosimum utile* (456), *Vochysia ferruginea* (373), *Peltogyne purpurea* (259), *Pouteria laevigata* (203), *Qualea polychroma* (208), *Aspidosperma spruceanum* (176), *Vantanea barbourii* (117), *Vochysia allenii* (144), *Tachigali versicolor*

Table 1: Diversity of plant species in one-hectare plots in the Golfo Dulce rainforest.

| Site in the Golfo Dulce region | d.b.h. | Number of indiv./ha | No. of spp./ha | Literature |
|-----------------------------------|---------|---------------------|----------------|-----------------------------------|
| Corcovado - Punta Llorona | ≥ 10 cm | 354 | 48 | HARTSHORN (1983) |
| Esquinas – ravine forest | ≥ 10 cm | 482 | 121 | HUBER (2005), WEISSENHOFER (2005) |
| Esquinas – slope forest (inland) | ≥ 10 cm | 527 | 133 | HUBER (2005), WEISSENHOFER (2005) |
| Esquinas – slope forest (coastal) | ≥ 10 cm | 588 | 108 | HUBER (2005), WEISSENHOFER (2005) |
| Esquinas –ridge forest | ≥ 10 cm | 847 | 179 | HUBER (2005), WEISSENHOFER (2005) |
| Bahia de Chal | ≥ 5 cm | 1161 | 201 | ZAMORA (unpubl.) |
| Agujas | ≥ 5 cm | 1098 | 189 | ZAMORA (unpubl.) |
| Corcovado- near Sirena | ≥ 5 cm | 881 | 149 | ZAMORA (unpubl.) |
| Esquinas - near ranger station | ≥ 5 cm | 871 | 188 | ZAMORA (unpubl.) |

(123), *Hyeronima alchomeoides* (118), *Ruptiliocarpon caracolito* (118), and *Vochysia megalophylla* (108).

Particular studies on the tree diversity revealed 98 species (≥ 10 cm d.b.h.) per hectare (HOLDRIDGE et al. 1971) in the Corcovado National Park, and 108 to 179 species (≥ 10 cm d.b.h.) per hectare in the Piedras Blancal National Park (Esquinas forest)(HUBER 2005, WEISSENHOFER 2005). An unpublished analysis of ZAMORA et al. shows that 149 to 201 species of woody plants (≥ 5 cm d.b.h.) per hectare are growing at four different sites in the rainforests around the Golfo Dulce rainforests (Table 1).

During the early evolutionary history of angiosperms, North and South America were separated. The isthmus between them formed in the Late Tertiary (c. 5 Million years ago). Though little is known about the of the geographical history of plants in Central and South America (CROAT & BUSEY 1975), it can be assumed that plants – like animals – migrated in both directions during the Late Tertiary.

Biogeography and floristic affinities

Central America belongs to the most diverse floristic regions in the world (GENTRY 1978) and forms part of the neotropical floristic kingdom (TAKHTAJAN 1986). About 32 angiosperm families are restricted to the neotropics (BORHIDI 1991), including Bromeliaceae, Cactaceae (except *Rhipsalis* in Africa and Madagascar), Caryocaraceae,

Cyclanthaceae, Heliconiaceae, Humiriaceae, Marcgraviaceae, Quinaceae, Theophrastaceae, Vochysiaceae, and others. Nonetheless, it is clear that the neotropical flora shares a common origin with the Palaeotropical flora and it may be assumed, at least for the flowering plants, that its roots are in the Palaeotropical kingdom. Many families have a pan-tropical distribution, e.g., Annonaceae, Bignoniaceae, Bombacaceae (now Malvaceae-Bombacoideae), Chrysobalanaceae, Clusiaceae, Lauraceae, Malpighiaceae, Moraceae, Myristicaceae, Proteaceae, Sapindaceae, Sapotaceae, and many others.

The neotropical plant kingdom can be subdivided into five floristic regions (TAKHTAJAN 1986). Costa Rica belongs to the Caribbean Region and is part of the Central American Province. This reaches from Mexico to the northern parts of South America, including the highly diverse region of the Chocó (Pacific lowlands of Colombia, Ecuador and northern Peru). A general comparison of the flora (vascular plants) of the ACOSA area (2.662 species counted so far) with the flora of Panama (CORREA et al. 2004), Chocó (FORERO & GENTRY 1989), Ecuador (JØRGENSEN et al. 1999), and Perú (BRAKO & ZARUCCHI 1993) shows that the percentage of shared species with those countries is: Panama: 79-80%, Chocó: 36%, Ecuador: 47%, and Peru: 35.6% (here, only seed plants have been considered).

A biogeographical analysis of the 312 tree species occurring in four 1 ha plots of the Esquinas forest (HU-

Table 2: Geographical distribution of seven selected plant families.

| | Life form | No. of species in the Golfo Dulce region | No. of species endemic to the Golfo Dulce region | No. of species. restricted to Central America | No. of species restricted to Central and South America | Neophytic |
|------------------|-----------|--|--|---|--|-----------|
| Chrysobalanaceae | tree | 22 | 6 | 3 | 12 | |
| Clusiaceae | tree | 30 | 2 | 8 | 20 | |
| Costaceae | herb | 15 | 3 | 2 | 9 | 1 |
| Lauraceae | tree | 55 | 10 | 23 | 22 | |
| Lecythidaceae | tree | 10 | 2 | 2 | 6 | |
| Marantaceae | herb | 25 | 8 | 4 | 13 | |
| Sapotaceae | tree | 40 | 4 | 11 | 24 | 1 |

BER 2005) showed that about half (161 species, 51%) are widespread in both South and Central America. More species have ranges extending to the east and south (to Panama: 251, Colombia: 197, Ecuador 166 and Amazon forest: 141) than to the west and north (Nicaragua: 186; Honduras; 109). The explanation is simple: the climatic situation is similar in north-western South America and the Golfo Dulce and leads one to expect a certain amount of floristic similarity. Apparently, these tree species originated in Amazonia, which is the main centre of tree diversity in the neotropics, and invaded Central America from there. This was enabled by the formation of the Central American land bridge and by the appropriate palaeoclimatic conditions for migration and establishment. In conclusion, the data clearly support the view that the Golfo Dulce area has strong floristic affinities to the forests of South America (GENTRY 1978, 1982a) and, in particular, to the Chocó region of north-western South America (STANDLEY 1937, HARTSHORN 1983, HARTSHORN & HAMMEL 1994, HUBER 1996, 2005).

Nonetheless, the high degree of endemism is remarkable. Of the 312 tree species analysed, 86 species are found exclusively in Central America. 99 species (c. 31%) occur only in the Pacific region (and not in the Caribbean region) of Costa Rica. This shows again that the Golfo Dulce forests reveal strong floristic differences to their counterparts on the Caribbean slope of Costa Rica. The reason for this is the presence of high mountain ranges separating the Pacific from the Caribbean slope. They are the result of strong tectonic movements in the region's eventful geological history.

In the frame of the present study, the geographical distribution of five selected tropical families of trees (Chrysobalanaceae, Clusiaceae, Lauraceae, Lecythidaceae, and Sapotaceae) and two families of herbs (Costaceae, Marantaceae) was analysed – with a total of 197 species (157 trees, 45 herbs). About 15% of the trees, and 27% of the herbaceous species proved endemic to southern Costa Rica (Table 2). In the Marantaceae, 8 out of 25 species (that is 32%) are endemic in the Golfo Dulce area.

Another (rough) analysis, relating to the whole flora of the ACOSA region (with 2.662 species counted so far) (ZAMORA, unpubl.), showed that about 150 species are endemic (c. 6%).

The considerable degree of endemism can be seen in context with the idea that the Golfo Dulce region was a natural refuge, cut off from the neighbouring forests during the Pleistocene (STILES 1983, HEPPNER 1991). Due to its isolated position, speciation occurred at an accelerated rate and many new species evolved in the fragmented forests. This resulted in “hot-spots” of biodiver-

Distritos ACOSA




Fig. 3: Map of the ACOSA region.



Fig. 4: *Costus osae* (Costaceae) is a herbaceous plant endemic to the Golfo Dulce region and is often found along small streams.



Fig. 5: *Calliandra grandifolia* (Fabaceae-Mimosoideae) is a beautiful tree endemic to the Golfo Dulce region.

sity, containing high numbers of endemic species. The MINAE (Ministerio de Natural, Ambiente y Energía de Costa Rica) qualifies the region around the Golfo Dulce as one of those areas of Costa Rica that are richest in endemic plants.

References

- ALLEN P.H. (1956): The rainforests of Golfo Dulce. — Stanford University Press, Stanford.
- BRAKO L. & J. ZARUCCHI (1993): Catalogue of the flowering plants and gymnosperms of Peru. — *Monogr. Syst. Bot. Missouri Bot. Gard.* **45**.
- BURGER W.C. (ed.) (1971 onw.): Flora Costaricensis. — *Fieldiana, Botany* **35**, **40**, and *Fieldiana, Botany New Series* **4**, **13**, **18**, **35**.
- BURGER W.C. (1980): Why are so many kinds of flowering plants in Costa Rica. — *Brenesia* **17**: 371-383.
- BURGER W. & C.M. TAYLOR (1993): Flora Costaricensis. — *Botany, New series*, No. **33**.
- CASTILLO M., FALLAS A & R. QUESADA (2007): Distribución y abundancia de árboles de dosel del bosque húmedo tropical en la Península de Osa. — Instituto Nacional de Biodiversidad-CTCBO-Critical Ecosystem Partnership Fund-Conservación Internacional, Instituto Tecnológico de Costa Rica, Informe Final, Cartago, C. R.
- CORREA M.D., GALDAMES C. & M.S. de STAFF (2004): Catálogo de las plantas vasculares de Panamá. Universidad de Panamá e Instituto Smithsonian de Investigaciones Tropicales, Panamá.
- CROAT T.B. & P. BUSEY (1975): Geographical affinities of the Barro Colorado island flora. — *Brittonia* **27**: 127-135.
- CONDIT R., PITTMANN N., LEIGH JR. E.G., CHAVE J., TERBORGH J., FOSTER R.B., NÚÑEZ P.V., AGUILAR S., VALENCIA R., VILLA G., MULLER-LANDAU H.C., LOSOS E. & S.P. HUBBELL (2002): Beta-diversity in tropical forest trees. — *Science* **295**: 666-669.
- DÍAZ R.E. (2008): Un ejemplo de intercambio científico entre Europa y América Latina: las investigaciones de la expedición científica Austriaca en Costa Rica (1930). — This volume.
- FORERO E. & A.H. GENTRY (1989): Lista Anotada de las Plantas del Departamento del Choco, Colombia. — Instituto de Ciencias Naturales-Museo de Historia Natural-Universidad Nacional de Colombia, Bogota.
- GENTRY A.H. (1978): Floristic knowledge and needs in Pacific Tropical America. — *Brittonia* **30**: 134-153.
- GENTRY A.H. (1982a): Phytogeographic patterns in northwest South America: a phytogeographical perspective. — In: HEDBERG I. (ed.), *Systematic botany, Plant utilization and biosphere conservation: symposium Uppsala*. Almqvist & Wiksell International, Stockholm.
- GENTRY A.H. (1982b): Phytogeographic patterns as Evidence for a Chocó refuge. — In: PRANCE G. (ed.), *Biological diversification in the tropics*. Colombia Univ. Press, Colombia.
- GENTRY A.H. (1986): Species richness and floristic composition of Choco region plant communities. — *Caldasia* **15**: 71-91.
- GÓMEZ L.D. (1986): Vegetación y clima de Costa Rica. Vol. 1. — EUNED, San José, Costa Rica.
- GOMEZ-LAURITO J. & L.A.O. FOURNIER (1985): Las familias y los generos de plantas lenosas de Costa Rica. — *Brenesia* **24**: 37-54.
- GRAYUM M.H., HAMMEL B.H., TROYO S. & N. ZAMORA (2004): Historia/History. — In: HAMMEL B.E., GRAYUM M.H., HERRERA C. & N. ZAMORA (eds), *Manual de plantas de Costa Rica. Vol. I. Introducción*. — *Monogr. Syst. Bot. Missouri Bot. Gard.* **97**: 1-50.
- HAMMEL B.E., GRAYUM M.H., HERRERA C. & N. ZAMORA (eds) (2003a): *Manual de plantas de Costa Rica. Vol II. Gimnospermas y Monocotiledóneas (Agavaceae-Musaceae)*. — *Monogr. Syst. Bot. Missouri Bot. Gard.* **92**: 1-694.
- HAMMEL B.E., GRAYUM M.H., HERRERA C. & N. ZAMORA (eds) (2003b): *Manual de plantas de Costa Rica. Vol III. Monocotiledóneas (Orchidaceae-Zingiberaceae)*. — *Monogr. Syst. Bot. Missouri Bot. Gard.* **93**: 1-884.
- HAMMEL B.E., GRAYUM M.H., HERRERA C. & N. ZAMORA (eds) (2004): *Manual de plantas de Costa Rica. Vol. I. Introducción*. — *Monogr. Syst. Bot. Missouri Bot. Gard.* **97**: 1-300.
- HAMMEL B.E., GRAYUM M.H., HERRERA C. & N. ZAMORA (eds) (2007): *Manual de plantas de Costa Rica. Vol VI. Dicotiledóneas (Haloragaceae-Phytolocaceae)*. — *Monogr. Syst. Bot. Missouri Bot. Gard.* **111**: 1-933.
- HARTSHORN G.S. (1983): Plants. Introduction. — In: JANZEN D.H. (ed.), *Costa Rican Natural History*, University of Chicago Press, Chicago.
- HARTSHORN G.S. & B.E. HAMMEL (1994): Vegetation types and floristic patterns. — In: MACDADE L.A., BAWA K.S., HESPENHEIDE H.A. & G.S. HARTSHORN (eds), *La Selva: ecology and natural history of a neotropical rainforest*, University of Chicago Press, Chicago.
- HEPPNER J.B. (1991): Faunal regions and the diversity of Lepidoptera. — *Trop. Lepidoptera* **2** (suppl. 1): 1-85.
- HERRERA W. (1985): Clima de Costa Rica. — In GÓMEZ L.D. (ed.), *Vegetación y clima de Costa Rica. Vol. 2*. EUNED. San José.
- HOLDRIDGE L.R., GENKE W.C., HATHWAG W.H., LIANG T. & J.A. TOSI JR. (1971): Forest environments in tropical life zones: A pilot study. — Pergamon Press, Oxford.
- HUBER W. (1996a): Floristische und biogeographische Untersuchungen in einem Tieflandregenwald in einer pazifischen Region in Costa Rica. — Unpubl. diploma thesis, University of Vienna, Vienna.
- HUBER W. (1996b): Untersuchungen zum Baumartenreichtum im "Regenwald der Österreicher" in Costa Rica. — *Carinthia II*, **186**: 95-106.
- HUBER W. (2005): Tree diversity and biogeography of four one-hectare plots in the lowland rainforest of the Piedras Blancas National Park ("Regenwald der Österreicher"), Costa Rica. — Unpubl. doctoral thesis, University of Vienna, Vienna.
- JORGENSEN P.M. & S. LEÓN YÁNEZ (1999): Catalogue of the vascular plants of Ecuador. — *Monogr. Syst. Bot. Missouri Bot. Gard.* **75**: 1-1181.
- PAMPERL S. (2001): Der Boden als Standortsfaktor eines baumartenreichen Tieflandregenwaldes in Costa Rica. — Vienna: Unpubl. diploma thesis, University of Vienna, Vienna.
- PITTMANN N.C.A., TERBORGH J.W., SILMAN M.R., NUNEZ P.V., NEILL D.A., CERON C.E., PALACIOS W.A. & M. AULESTIA (2002): A comparison of tree species diversity in two upper Amazonian forests. — *Ecology* **83**: 3210-3224.
- PRANCE G.T. (1994): A comparison of the efficacy of higher taxa and species numbers in the assessment of biodiversity in the neotropics. — *Philosophical Transactions of the Royal Society of London* **345**: 89-99.

- QUESADA F.J., JIMÉNEZ Q., ZAMORA N., AGUILAR R. & J. GONZÁLEZ (1997): Arboles de Peninsulæ de Osa. — Heredia: INBio.
- SISTEMA NACIONAL DE AREAS DE CONSERVACIÓN (SINAC) DEL MINISTERIO DEL AMBIENTE Y ENERGÍA (MINAE). (2007): guas 11: Propuesta de Ordenamiento Territorial para la Conservación de la Biodiversidad de Costa Rica. Volumen 1: Análisis de Vácíos en la Representatividad e Integridad de la Biodiversidad Terrestre, San Jose, C.R.
- STANDLEY P.C. (1937): Flora of Costa Rica. — Publ. Field Mus. Nat. Hist., Bot. Ser. **18**: (1)-(4).
- STILES F.G. (1983): Birds. Introduction. — In JANZEN D.H. (ed.), Costa Rican natural history. University of Chicago Press, Chicago: 502-530.
- TAKHTAJAN A. (1986): Floristic regions of the World. — Univ. Calif. Press, Berkeley.
- VAUGHAN C.S. (1981): Parque Nacional Corcovado: plan de manejo y desarrollo. — Heredia: Universidad Nacional.
- WEBER A. (2008): Otto Porsch and the scientific goals and results of the Austrian Costa Rica expedition 1930. — This volume.
- WEBER A., HUBER W., WEISSENHOFER A., ZAMORA N. & G. ZIMMERMANN (2001): An introductory field guide to the flowering plants of the Golfo Dulce rainforests, Costa Rica. — *Stapfia* **78**: 1-462.
- WEISSENHOFER A. (1996): Ökologie und Struktur eines Tieflandregenwaldes in der Pazifikregion Costa Ricas. — Unpubl. diploma thesis, University of Vienna, Vienna.
- WEISSENHOFER A. (1997): Untersuchungen zur Ökologie und Struktur im "Regenwald der Österreicher" in Costa Rica. — *Carinthia II* **187**: 67-80.
- WEISSENHOFER A. (2005): Structure and vegetations dynamics of four selected one hectare forest plots in the lowland rain forests of the Piedras Blancas National Park ("Regenwald der Österreicher"), Costa Rica, with notes on the vegetation diversity of the Golfo Dulce region. — Unpubl. doctoral thesis, University of Vienna, Vienna.
- WEISSENHOFER A., HUBER W., ZAMORA N., WEBER A. & J. GONZÁLEZ (2001): A brief outline of the flora and vegetation of the Golfo Dulce region. — *Stapfia* **78**: 11-14.
- WEISSENHOFER A. & W. HUBER (2008): The climate of the Esquinas rainforest. — In this volume.
- ZAMORA N., HAMMEL B.E. & M.H. GRAYUM (2004): Vegetación/Vegetation. — In: HAMMEL B.E., GRAYUM M.H., HERRERA C. & N. ZAMORA (eds), *Manual de Plantas de Costa Rica*. Vol. I. Introducción, *Monogr. Syst. Bot. Missouri Bot.* **97**: 91-216.

Addresses of authors:

Werner HUBER
 Anton WEISSENHOFER
 Anton WEBER
 Department of Palynology and Structural Botany
 Faculty Centre of Botany
 University Vienna
 Rennweg 14
 A-1030 Vienna, Austria
 E-mail: werner.huber@univie.ac.at
 anton.weissenhofer@univie.ac.at
 anton.weber@univie.ac.at

Nelson ZAMORA
 INBio, Instituto Nacional de Biodiversidad
 Santo Domingo de Heredia, Costa Rica
 E-mail: nzamora@inbio.ac.cr

Contents/Indice

Prefaces and introduction — Prólogos y introducción

| | |
|---|----|
| Dr. Pedro LEON Coordinator, President Aria's Initiative on Peace with Nature | 11 |
| O. Univ.-Prof. Dr. Georg WINKLER Rector of the University Vienna | 12 |
| Dr. Josef PÜHRINGER Governor of Upper Austria | 13 |
| Introduction Introducción | 14 |

ABIOTIC ASPECTS — FACTORES ABIÓTICOS

Geography — Geografía

| | |
|--|----|
| Geography of the Golfo Dulce region (survey) Geografía de la región del Golfo Dulce (sinopsis) WEISSENHOFER A., HUBER W. & KLINGLER M. | 19 |
|--|----|

Geology — Geología

| | |
|--|----|
| Outline of the geology of the Golfo Dulce Region (Costa Rica) and its surroundings in Central America (survey) Vista de conjunto de la geología de la Región del Golfo Dulce (Costa Rica) y de sus inmediaciones en América Central (sinopsis) MALZER O. & FIEBIG M. | 23 |
| Geological and mineralogical investigations of the lithologies and their weathering products in a study area south-west of the field station "La Gamba", Golfo Dulce, Costa Rica Investigación geológica y mineralógica de las rocas y sus productos de meteorización, en un área al suroeste de la estación "La Gamba", Golfo Dulce, Costa Rica SCHEUCHER L.E.A., VORTISCH W. & LAGUNA-MORALES J. | 31 |
| Anthropogenic and natural radionuclides in soil of a tropical rainforest of Southern Costa Rica Radionúclidos antropogénicos y naturales en el suelo de un bosque lluvioso tropical del sur de Costa Rica BOSSEW P., HUBMER A. & STREBL F. | 47 |

Climate — Clima

| | |
|---|----|
| The climate of the Esquinas rainforest (survey) El clima del bosque lluvioso Esquinas (sinopsis) WEISSENHOFER A. & HUBER W. | 59 |
|---|----|

PLANT BIOLOGY — BIOLOGÍA DE PLANTAS

Ecosystems and vegetation — Ecosistemas y vegetación

Ecosystem diversity in the Piedras Blancas National Park and adjacent areas (Costa Rica), with the first vegetation map of the area

Diversidad de ecosistemas en el Parque Nacional Piedras Blancas y áreas adyacentes (Costa Rica), con la primera presentación de una mapa vegetacional

WEISSENHOFER A., HUBER W., KOUKAL T., IMMITZER M., SCHEMBERA E., SONTAG S., ZAMORA N. & WEBER A. 65

Plant diversity — Diversidad de plantas

Plant diversity and biogeography of the Golfo Dulce region, Costa Rica (survey)

Diversidad vegetal y biogeografía de la región de Golfo Dulce, Costa Rica (sinopsis)

HUBER W., WEISSENHOFER A., ZAMORA N. & WEBER A. 97

Alien plants and invasion patterns in different habitats of the Golfo Dulce area, Costa Rica

Plantas exóticas y patrones de invasión en diferentes hábitat del área de Golfo Dulce, Costa Rica

HUBER W., WEISSENHOFER A. & ESSL F. 105

Survey of Rubiaceae in the Golfo Dulce area, Costa Rica:
New species, combinations and name changes since 2001

Investigaciones en Rubiaceae en el área de Golfo Dulce, Costa Rica:
nuevas especies, combinaciones y cambios de nombre desde 2001

WILL S. & KIEHN M. 111

Medicinal plants in La Gamba and in the Esquinas rain forest

Plantas medicinales en La Gamba y de la selva tropical Esquinas

LÄNGER R. 121

Life forms — Formas de vida vegetal

Plant life forms in the Golfo Dulce region and other neotropical rainforests (survey)

Formas de vida vegetal en la región de Golfo Dulce y en otros bosques lluviosos neotropicales (sinopsis)

HIEZ P. 129

Terrestrial litter trappers in the Golfo Dulce region: diversity, architecture and ecology of a poorly known group of plant specialists

Plantas captadoras de hojarasca en la región de Golfo Dulce: diversidad, arquitectura y ecología de un grupo de plantas especialistas poco conocido

WEISSENHOFER A., HUBER W., WANEK W. & WEBER A. 143

Ecophysiology — Ecofisiología

Primary production and nutrient cycling in lowland rainforests of the Golfo Dulce region

Producción primaria y ciclo de nutrientes en bosques lluviosos de tierras bajas de la región de Golfo Dulce

WANEK W., DRAGE S., HINKO N., HOFHANSL F., PÖLZ E.-M., RATZER A. & RICHTER A. 155

Fungi and lichens — Hongos y líquenes

Diversity and ecology of fungi in the Golfo Dulce region (survey)

Diversidad y ecología de hongos en la región del Golfo Dulce (sinopsis)

PIEPENBRING M. & RUIZ-BOYER A.

179

The lichens of the Golfo Dulce region (survey)

Líquenes de la región de Golfo Dulce (sinopsis)

BREUSS O.

193

ANIMAL BIOLOGY — BIOLOGÍA DE ANIMALES

Spiders — Arañas

Cupiennius (Araneae, Ctenidae): Biology and sensory ecology of a model Spider

Cupiennius (Araneae, Ctenidae): Biología y ecología sensorica de una araña modelo

BARTH F.G.

211

Key to the genus *Cupiennius* (Araneae, Ctenidae)

Clave de determinación de genus *Cupiennius* (Araneae, Ctenidae)

BARTH F.G. & CORDES D.

225

Insects — Insectos

Diversity, biogeography and ecology of insects in the Pacific lowlands of Costa Rica, with emphasis on La Gamba (survey)

Diversidad, biogeografía y ecología de los insectos en las tierras bajas del Pacífico de Costa Rica con énfasis en La Gamba (sinopsis)

SCHULZE C.H.

229

Banderillas: Effects of deforestation on dragonflies (Insecta, Odonata) in the Pacific lowland of Costa Rica

Banderillas: Efectos de la deforestación sobre libélulas (Insecta, Odonata) en la tierra baja Pacífica en Costa Rica

HOFHANSL F.P. & SCHNEEWEIHS S.

237

Longhorn beetles (Coleoptera, Cerambycidae) of the Golfo Dulce region, Costa Rica

Cerambícidos (Coleoptera, Cerambycidae) de la región de Golfo Dulce, Costa Rica

HUBWEBER L.

249

Diversity of Euglossini (Hymenoptera, Apidae) in primary and secondary lowland rainforests in south-western Costa Rica

Diversidad de Euglossini (Hymenoptera, Apidae) en bosques lluviosos de tierras bajas primarios y secundarios en el sudoeste de Costa Rica

GRUBER M.H., MORAWETZ L. & WIEMERS M.

257

Stingless bees of the Golfo Dulce region, Costa Rica (Hymenoptera, Apidae, Apinae, Meliponini)

Las abejas sin aguijón de la región de Golfo Dulce, Costa Rica (Hymenoptera, Apidae, Apinae, Meliponini)

JARAU S. & BARTH F.G.

267

Butterfly diversity of the Piedras Blancas National Park and its vicinity –
a preliminary assessment (Lepidoptera: Papilionidae & Hesperioidea)

Diversidad de mariposas del Parque Nacional Piedras Blancas y zonas cercanas –
una evaluación preliminar (Lepidoptera: Papilionidae & Hesperioidea)

WIEMERS M. & FIEDLER K.

277

Feeding behaviours of neotropical butterflies (Lepidoptera, Papilionoidea)

Ingestión de alimentos en mariposas neotropicales (Lepidoptera, Papilionoidea)

KRENN H.W.

295

Amphibians and reptiles — Anfibios y reptiles

The amphibians and reptiles of the Golfo Dulce region (survey)

Los anfibios y reptiles de la región del Golfo Dulce (sinopsis)

HÖBEL G.

305

Plasticity and geographic variation in the reproductive ecology of gladiator frogs,
particularly *Hypsiboas rosenbergi*

Plasticidad y variación geográfica en la ecología reproductiva de ranas gladiadoras, especialmente *Hypsiboas rosenbergi*

HÖBEL G.

329

Reproductive behaviour of the glass frog *Hyalinobatrachium valerioi*
(Anura: Centrolenidae) at the tropical stream Quebrada Negra (La Gamba, Costa Rica)

Comportamiento reproductivo de la rana de cristal *Hyalinobatrachium valerioi* (Anura: Centrolenidae)
en el arroyo tropical Quebrada Negra (La Gamba, Costa Rica)

VOCKENHUBER E.A., HÖDL W. & KARPEN U.

335

Birds — Aves

Birds of La Gamba – a call for research and scientific collaboration

Aves de La Gamba – un llamado para investigación y colaboración científica

AUBRECHT G. & SCHULZE C.H.

349

The birds of La Gamba (survey)

Los pajaros de La Gamba (sinopsis)

TEBB G.

353

Habia atrimaxillaris (Dwight & Griscom) 1924 – the black-cheeked ant-tanager.
History of an endemic bird species from SW Costa Rica, from discovery to endangered status

Habia atrimaxillaris (Dwight & Griscom) 1924 – tangara hormiguera cabecinegra.

Historia de una especie de ave endémica del Sudoeste de Costa Rica – desde su descubrimiento a su estatus de peligro

AUBRECHT G.

381

Bird assemblages of forested and human-modified countryside habitats
in the Pacific lowlands of southern Costa Rica

Grupos de aves de hábitat boscosos y rurales en las tierras bajas del Pacífico del sur de Costa Rica

SCHULZE C.H. & RIEDL I.

395

Mammals — Mamíferos

Mammals of the Piedras Blancas National Park, Costa Rica: species composition, habitat associations and efficiency of research methods – a preliminary overview (survey)

Mamíferos del Parque Nacional Piedras Blancas, Costa Rica: composición de especies, asociaciones de hábitat y eficiencia de métodos de investigación – una panorámica preliminar (sinopsis)

LANDMANN A., WALDER C., VORAUER A. & EMSER T.

409

Bats of the La Gamba region, Esquinas rain forest, Costa Rica: species diversity, guild structure and niche segregation

Murciélagos de la región de La Gamba, bosque lluvioso Esquinas, Costa Rica: diversidad específica, estructura gremial y segregación de nichos

LANDMANN A., WALDER C., VORAUER A., BOHN S. & WEINBEER M.

423

Limnology — Limnología

The river network of the Piedras Blancas National Park, Costa Rica (survey)

La red fluvial del Parque Nacional Piedras Blancas, Costa Rica (sinopsis)

TSCHELAUT J., PICHLER C., WEISSENHOFER A. & SCHIEMER F.

441

Macroinvertebrates and leaf litter decomposition in a neotropical lowland stream, Quebrada Negra, Costa Rica

Macroinvertebrados y descomposición de residuos de hojas en un curso de agua de tierras bajas neotropical, Quebrada Negra, Costa Rica

TSCHELAUT J., WEISSENHOFER A. & SCHIEMER F.

457

The role of leaf anatomy and tannins in litter decay in a tropical stream

El rol de la anatomía foliar y de los taninos en la descomposición de la hojarasca en un arroyo tropical

RIEMERTH A., GUSENLEITNER M. & SCHIEMER F.

467

Freshwater bryozoans in La Gamba (Costa Rica: Piedras Blancas National Park): a general introduction

Briozoos de agua dulce en La Gamba (Costa Rica: Parque Nacional Piedras Blancas): una introducción

WÖSS E.R.

485

Ecology of fishes of Quebrada Negra, Costa Rica, a first order neotropical lowland stream

Ecología de los peces de Quebrada Negra, Costa Rica, río neotropical de primer orden de tierras bajas

PICHLER C. & SCHIEMER F.

495

PLANT-ANIMAL INTERACTIONS — INTERACCIONES DE PLANTAS Y ANIMALES

Pollination — Polinización

Pollination in the plants of the Golfo Dulce area (survey)

Polinización en las plantas del área de Golfo Dulce (sinopsis)

WEBER A.

509

Phenology and pollination of *Ceiba pentandra* (Bombacaceae) in the wet forest of south-eastern Costa Rica

Fenología y biología de la polinización de *Ceiba pentandra* (Bombacaceae) en el bosque húmedo del sudeste de Costa Rica

ROJAS-SANDOVAL J., BUDDE K., FERNÁNDEZ M., CHACÓN E., QUESADA M. & LOBO J.A.

539

Phenology of tree species of the Osa Peninsula and Golfo Dulce region, Costa Rica

Fenología de especies de árboles de la Península de Osa y la región de Golfo Dulce, Costa Rica

LOBO J., AGUILAR R., CHACÓN E. & FUCHS E.

547

Style release experiments in four species of Marantaceae from the Golfo Dulce area, Costa Rica

Experimentos acerca de la sensibilidad del estilo en cuatro especies de marantáceas del área del Golfo Dulce, Costa Rica

CLASSEN-BOCKHOFF R. & HELLER A.

557

Notes on the pollination of the perfume flowers of *Gloxinia perennis* (Gesneriaceae) by euglossine bees

Notas sobre la polinización de las flores perfumadas de *Gloxinia perennis* (Gesneriaceae) por abejas euglossine

WITSCHNIG G., HICKL C. & WEBER A.

573

Scientific work of Austrian students in the “Austrian rainforest” (Piedras Blancas National Park, Costa Rica), with special regard to pollination studies

Trabajo científico de los estudiantes en el “Bosque lluvioso de los Austriacos”

(Parque Nacional Piedras Blancas, Costa Rica), con especial consideración en los estudios de polinización

WEBER A.

579

Ants and plants — Hormigas y plantas

A house in the tropics: full pension for ants in *Piper* plants

Una casa en el tropico: pensión completa para hormigas en plantas de *Piper*

FISCHER R. & MAYER V.

589

Does nectar production reduce herbivore pressure on *Passiflora* species (Passifloraceae) in a tropical rainforest in Costa Rica?

Puede la producción de néctar reducir la presión de herbivoría en especies de *Passiflora* (Passifloraceae) en un bosque tropical de Costa Rica?

THURNER M. & MAYER V.

599

HUMAN ASPECTS — ASPECTOS HUMANOS

History and development — Historia y desarrollo

Indigenous societies of the south east of Costa Rica, 15th century

Sociedades indígenas del sudeste de Costa Rica, siglo XVI

BARRANTES CARTÍN C.

609

The stone balls of Palmar

Las esferas de piedra de Palmar

STEPHENS C.

631

| | |
|---|-----|
| Impacts of the United Fruit Company in Southwest Costa Rica Impacto de la United Fruit Company en el Suroeste de Costa Rica STEPHENS C. | 635 |
|---|-----|

| | |
|--|-----|
| Corcovado National Park – almost a banana plantation El Parque Nacional Corcovado – casi una plantación bananera STEPHENS C. | 645 |
|--|-----|

| | |
|---|-----|
| Case study: economic and structural settlement changes and their consequences in the community of La Gamba, Golfo Dulce region Estudio de caso: Cambios estructurales y económicos de la población y sus consecuencias en la comunidad La Gamba, región del Golfo Dulce KLINGLER M. | 649 |
|---|-----|

History of nature exploration — Historia de exploración de la naturaleza

| | |
|---|-----|
| Un ejemplo de intercambio científico entre Europa y América Latina: las investigaciones de la expedición científica Austríaca en Costa Rica (1930) An example of scientific interchange between Europe and Latin America: The investigations of the Austrian scientific expedition to Costa Rica (1930) DÍAZ BOLAÑOS R.E. | 657 |
|---|-----|

| | |
|--|-----|
| Otto Porsch and the scientific goals and results of the Austrian Costa Rica expedition 1930 Otto Porsch y los objetivos y resultados científicos de la expedición Austriaco-Costarricense de 1930 WEBER A. | 667 |
|--|-----|

History of nature conservation — Historia de protección de la naturaleza

| | |
|---|-----|
| Before Corcovado: Early conservation initiatives on the Osa Peninsula (survey) Antes del Corcovado: Primeras iniciativas pro-conservación en la Península de Osa (sinopsis) CHRISTEN C.A. | 675 |
|---|-----|

| | |
|--|-----|
| Looking back to the foundation of the Corcovado National Park, a crown jewel of nature in Costa Rica Una mirada retrospectiva a la fundación del Parque Nacional Corcovado, una corona de joyas de la naturaleza en Costa Rica UGALDE A. | 683 |
|--|-----|

| | |
|--|-----|
| In defence of local livelihoods, the forest and the Golfo Dulce: the campaign against “Ston Forestal” in the 1990s and its historical roots En defens de los medios de vida locales, el bosque y el Golfo Dulce: la campaña contra Ston Forestal en los 1990 y sus raíces históricas VAN DEN HOMBERGH H. | 693 |
|--|-----|

Current conservation projects — Proyectos actuales de protección de la naturaleza

| | |
|--|-----|
| The Osa biological corridor in the context of the mesoamerican biological corridor El corredor biológico Osa en el contexto del corredor biológico mesoamericano GARCÍA R. | 701 |
|--|-----|

Conectividad entre el Parque Nacional Piedras Blancas y la Fila de Cal

The link between the Piedras Blancas National Park and the Fila de Cal

MORERA C. & ROMERO M.

707

The Biological Corridor Project in the Piedras Blancas National Park, Costa Rica.

A project to preserve the biodiversity by reforestation and alternative culture, with support of the community La Gamba by new marketing strategies

El proyecto de corredor biológico en el Parque Nacional Piedras Blancas, Costa Rica.

Un proyecto para preservar la biodiversidad mediante la reforestación y cultivos alternativos, con el apoyo de la comunidad de la Gamba de nuevas estrategias de comercialización

WEISSENHOFER A., BARQUERO M., HUBER W., MAYER V. & NÁJERA UMAÑA J.

715

Reintroducción de la Lapa Roja (*Ara macao*) en Playa San Josecito, Golfito

Reintroduction of the scarlet macaw (*Ara macao*) to Playa San Josecito, Golfito

VARELA BENAVIDES I. & JANIK D.

725

The project "Rainforest of the Austrians"

El proyecto "Bosque de los Austriacos"

SCHNITZLER M.

733

The "Tropical Research Station La Gamba" – science, education and nature conservation in Costa Rica

La "Estación Tropical La Gamba" – ciencia, educación y conservación en Costa Rica

ALBERT R. & WEBER A.

739

Ecotourism and local development — Ecoturismo y desarrollo local

Ecotourism in La Gamba. An economical and ecological alternative for the residents of La Gamba?

Ecoturismo en La Gamba. Una alternativa económica y ecológica para los residentes de La Gamba?

FAHRNBERGER M.

743

Sinergias entre ecoturismo y desarrollo local en la península de Osa, Costa Rica

Synergistic effects of ecotourism and local development on the Osa Peninsula, Costa Rica

MORERA C.

755

APPENDIX — APÉNDICE

Authors' addresses

Direcciones de los autores

763

Vegetation map of the Piedras Blancas Nationalpark, Golfito Forest Reserve and adjacent areas

Mapa de vegetación del Parque Nacional Piedras Blancas, Reserva Forestal Golfito y áreas adyacentes